

Superconducting String Theory (Gravity Explanation)

Sergio P. F.¹

¹Independent Researcher, graduated from Computer Science, Madrid, Spain

Correspondence: Sergio P. F., Independent Researcher, graduated from Computer Science, Madrid, Spain. E-mail: sergiopf@gmail.com

Received: January 12, 2017

Accepted: January 22, 2017

Online Published: January 31, 2017

doi:10.5539/apr.v9n1p99

URL: <http://dx.doi.org/10.5539/apr.v9n1p99>

Abstract

Gravity explained by a new theory, 'Superconducting String Theory (SST)', completely opposite from current field emission based and inspired on originals string theories. Strengths are decomposed to make strings behave as one-dimensional structure with universe acting as a superconductor where resistance is near 0 and the matter moves inside. Strong nuclear force, with an attraction of 10.000 Newtons is which makes space to curve, generating acceleration, more matter more acceleration. Electromagnetic moves in 8 decimals, gravity is moved to more than 30 decimals to work as a superconductor.

Keywords: superconducting string theory, string theory, strong nuclear force, gluon, unified field theory, gravity, relativity

1. Introduction

The 'Theory of Everything' is a hypothetical theory of physics that explains and connects all known physical phenomena into one. There is a possible solution to the origin of gravity force, postulating it as angular piece of this theory, this solution erases gravity as one of the fundamental forces of nature and unifies it with strong nuclear force.

Let's analyze the forces that occur in the universe transforming string theory. It allows to explain many physical behaviors that without its existence would be practically impossible to understand, even so, these strings have not been able to be discovered and are only that, a theory that serves as an important support to the world of physics. One of the best known theoretical applications about them is how their vibration can provoke the creation of matter, but this is not about theories already written, we are going to place these strings in a simpler way to answer some doubts in subatomic world.

This theory uses 4 dimensions in space and a behavior as one dimension in strings with superconducting capacities. Like an elastic band between V-shaped sticks where the elastic band slides down, the strong nuclear force, forces these strings to bend to fall dawn.




Figure 1. String example

It's not directly related to electromagnetism.

2. Actors

2.1 String Theory

String theory is a theoretical framework in which the point-like particles of particle physics are replaced by one-dimensional objects called strings. Each string that we cross would be the minimum distance that can be traversed during a displacement.

We can note two important qualities of strings:

- Distance to the most distant object detected by the human being is more than 30 billion light years, that means there are beams of light which are able to travel that distance without decreasing its speed (they modify only its wavelength). Like light, an object can move into space for a practically unlimited period, as long as it doesn't find a force to stop it. If strings exist, they act as a superconductor of matter with a resistance near 0.
- In order to generate waves it's easier into a strongly linked structure. Gravitational waves behave like ocean waves which are similar to an uptight net, these tensions can be decomposed as one-dimensional structure for its study. Strings, at same time, could be one or zero-dimensional, like points under extreme bound forces, think about them as something tenser than any cable that holds the heaviest bridge in the world.

The new framework we have drawn would be a set of extremely tense strings, with a practically infinite matter conduction capacity.



Figure 2. Example of quantum entanglement. Gravity waves behave as an uptight net

2.2 Strong Nuclear Force

Strong nuclear force is another variable. This force allows the atomic nucleus to remain together, being the strongest of the so-called fundamental interactions (gravitational, electromagnetic, strong nuclear, and weak nuclear). Gluon is in charge of this interaction, it has a scope not greater than 10 to the power of -15 meters, preventing matter to separate by a constant attraction force between quarks of maximum 10.000 N (F).




Figure 3. Gluon into vacuum (Special Research Centre for the Subatomic Structure of Matter)

This real picture illustrates the three-dimensional structure of gluon-field configurations, describing the vacuum properties. The volume of the box is 2,4 by 2,4 by 3,6 fm. Contrary to the concept of an empty vacuum, this

induces chromo-electric and chromo-magnetic fields in its lowest energy state. The frame rate into this example is billions of billions frames per second (FPS).

3. Superconducting String Theory (SST)

3.1 Fundamentals

We have created a scenario with a superconductor of matter interacting with a force that makes that matter hold together, but, how can they interact? The most simple is to think about two V-shaped sticks (simulating the strings), and an elastic band that tight them at the most opened side (it would simulate the gluon, with size 10 to the power of -15 meters). If sticks are sufficiently lubricated and tense, what does the elastic band do? It will slide to the thinnest side. More elastic bands, more force will be exerted on the sticks to join them, so next bands will slide even faster (equally, more mass causes more attraction).

We are talking about unknown limits in known world, such as infinite conduction or tensions never seen in materials.




Figure 4. Implied forces

Suddenly, we have erased one of the fundamental forces of nature, gravity force doesn't really exist, exists the strong nuclear force interacting with strings. I have called this theory 'Superconducting String Theory (SST)'.

3.2 Calculations

Apply formulas from inclined planes (Newton's second law). Simulation is in horizontal direction.

Friction is imperceptible and acceleration down the plane is matched with gravity acceleration in our planet. Vertical force is not gravity force, it is gluon force. It can be considered vertical angle, but is depreciable and gluon force is estimated, so we keep 10.000 N.




Figure 5. Inclined plane forces (gluon catching strings)

$$m_1 \times g \times \text{sen}(x) = m_2 \times a$$

Match vertical force (F_1) with the attraction force from one gluon (estimated max. constant 10.000 N).

$$F_1 = m_1 \times g = 10.000 \text{ N} \tag{1.1}$$

$$F_1 \times \text{sen}(x) = m_2 \times a$$

Convert variables to metric system considering the mass of the gluon ($0,0002 \text{ eV}/c^2$).

$$1\text{eV}/c^2 = 1,782662 \times 10^{-36} \text{ kg} \tag{1.2}$$

$$m_2 = 0,0002 \times 1,782662 \times 10^{-36} = 3,565324 \times 10^{-40} \text{ kg}$$

$$a = 9,8 \text{ m/s}^2$$

$$F_2 = m_2 \times a = 3,565324 \times 10^{-40} \times 9,8 = 3,49401752 \times 10^{-39} \text{ N}$$

Angle calculation of all strings under gluon influence.

$$x = \text{arcsen}(F_2 / F_1) \tag{1.3}$$

$$F_2 / F_1 = 3,49401752 \times 10^{-43} \text{ N}$$

$$x = \text{arcsen}(3,49401752 \times 10^{-43}) = 3,49401752 \times 10^{-43} \text{ }^\circ$$




Figure 6. θ angle of all strings gluon can catch

4. Conclusions

It can explain:

- What is gravity force.
- Unified field theory between gravity and strong nuclear force.
- New behavior in dark matter because of differences in density of the superconductor. Lower density generates bigger angle, this implies bigger attraction force. If threads are separated, matter (m_2) becomes energy (F_1). Some places at universe could have bigger accelerations because of this effect; this means much less dark matter (which is stimated at approximately 27% of the mass and energy in the observable universe).




Figure 7. Increase the separation, increase the force exerted. Matter is transformed into energy

$$F_1 \times \text{sen}(x) = m_2 \times a \tag{2.1}$$

$$1\text{eV}/c^2 = 1,782662 \times 10^{-36} \text{ kg}$$

$$m_2 = 0,0002 \times 1,782662 \times 10^{-36} = 3,565324 \times 10^{-40} \text{ kg}$$

Calculate acceleration in relation to angle.

$$F_2 = m_2 \times a = 3,565324 \times 10^{-40} \times a \text{ kg/(m/s}^2) \tag{2.2}$$

$$x = \arcsen(F_2 / F_1)$$

$$F_2 / F_1 = (3,565324 \times 10^{-40} \times a) / 10.000 = (3,565324 \times 10^{-44} \times a)$$

$$x = \arcsen(3,565324 \times 10^{-44} \times a) = (3,565324 \times 10^{-44} \times a)^\circ$$

Bigger angle generates more acceleration.

$$a = (x / 3,565324 \times 10^{-44}) \text{ m/sg}^2 \tag{2.3}$$

- Einstein field equations. Apply Hooke's law to gluon force as a vertical spring to calculate tensions.




Figure 8. Energy don't disappear becomes matter

Gluon force has negative correlation when transforming into matter, decrease F1 or increase m2 implies less acceleration.

$$a = F_1 \times \text{sen}(x) / m_2 \tag{3.1}$$

$$a = 5.000 \times \text{sen}(3,49401752 \times 10^{-43}) / 3,565324 \times 10^{-40}$$

$$a = 4,9 \text{ m/s}^2$$

- Spherical and circular movements at planets and galaxies. Space deforms not proportionally to create more acceleration near accumulation of matter, behaving like an elastic material; this behavior can be quantified by Young's modulus which represents the factor of proportionality in Hooke's law at non-linear systems.




Figure 9. Angle originated by big amounts of matter has an interaction over very large distances

The elastic modulus or Young's modulus (E) depends on the force exerted by matter (σ) and the deformation at each point of the resulting vector (ε).

$$E = \Delta\sigma / \Delta\epsilon$$

Force exerted by the angle ($F\theta$), increases (Δ) faster than force exerted by gluon (F_1) and its relation with matter (m_2).

$$\Delta F\theta > \Delta F_1 / \Delta m_2 \tag{4.1}$$

This gluon-matter relation also modifies the density of the superconductor in space, since it induces their approach, therefore we can speak of the existence of a bulk modulus (K).

$$K = -V (\Delta p / \Delta V)$$

The bulk modulus (K) depends on pressure changes (p) and volume (V).

Other properties as volume viscosity also called bulk viscosity can be applied.




Figure 10. All these variables help to create shapes in galaxies like the golden spiral ($\phi = 1,6180$)

- Gravitational constant ($G = 6,67408 \times 10^{-11} \text{ m}^3\text{kg}^{-1}\text{s}^{-2}$) and its problem to measure with high accuracy since it can be related to the density exposed.
- Schrödinger equation, to describe how the quantum state of a quantum system changes with time, similar to Newton's second law.
- Planck's length ($1,616229 \times 10^{-35} \text{ m}$) which can indicate the distance between strings.
- Gluon size and its larger size far from earth.
- Black holes.

Rereferences

Newton, I. (1687). *Philosophiae naturalis principia mathematica*.
 Hooke, R. (1678). *The extension is proportional to the force*.
 Euler, L. (1768). *Institutionum calculi integralis* (Vol. 1). imp. Acad. imp. Saent.

Copyrights

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.
 This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/4.0/>).